

	Before Propensity score matched		Propensity score matched		Effect Measure	
	HA n = 40,764	SA n = 191,391	HA n = 40,760	SA n = 40,760		
Categorical value					RD (95% CI)	RR (95% CI)
Reoperation (count (%))	517 (1.27)	1,565 (0.82)	517 (1.27)	380 (0.93)	0.34 (0.19 to 0.48)	1.36 (1.19 to 1.55)
Percutaneous abscess drainage (count (%))	187 (0.46)	964 (0.50)	187 (0.46)	213 (0.52)	-0.06 (-0.16 to 0.03)	0.88 (0.72 to 1.07)
Endoscopic intestinal dilation (count (%))	29 (0.07)	66 (0.03)	29 (0.07)	17 (0.04)	0.03 (0.00 to 0.06)	1.71 (0.94 to 3.10)
Endoscopic intestinal hemostasis (count (%))	17 (0.04)	343 (0.18)	17 (0.04)	80 (0.20)	-0.16 (-0.20 to -0.11)	0.21 (0.13 to 0.36)
Continuous value					Difference (95% CI)	
Mean general anesthesia time (min. (SD))	273.51 (88.81)	277.89 (93.96)	273.51 (88.81)	272.02 (93.19)	1.49 (0.25 to 2.74)	
Mean length of postoperative stay (day (SD))	15.58 (14.82)	15.18 (13.75)	15.58 (14.82)	15.79 (14.08)	-0.21 (-0.40 to -0.01)	